

Diabetes Medication Administration Form [Part A]

DUE: June 1st. Forms submitted after June 1st may delay processing for new school year.

Provider Medication Order Form | School Year 2023-24 Please fax all DMAFs to 347-396-8932/8945

Student Last Name: First N		First Na	rst Name:		Date of Birth:	☐ Male ☐ Female	☐ Male OSIS #				
School ATSDBN / Name: Address:				Borough:		DOE District:	Grade:	Class:			
HEALTH CARE PRACTITIONER COMPLETES BELOW [Ple						ase see 'Provider Guidelii	nes for DMAF Com	pletion']			
☐ Type 1 Diabetes					Recent A1c						
Other Diagnosis:						Date		Resu	ılt%		
Orders written will b	e implemented	when subm	nitted an	d approved. If you wis	sh to delay or	ders for September 2023 pl	ease check here				
					RGENCY O	RDERS					
		ere Hypoglyc		044	Risk for Ketones or Diabetic Ketoacidosis (DKA) 11 □ Test ketones if bG > mg/dl or if vomiting, or fever > 100.5 F						
Glucagon	GVOKE	Glucagon ar Bagsin		Zegalogue	l or						
□ 1 mg	□ 1 mg	☐ 3 mg		□ 0.6 mg SC □ Test ketones if bG > mg/dl for the 2nd time that day (at least 2 hrs. apa					2 hrs. apart), or if		
□mg	□ mg	Intranasa		ay repeat in 15 min if	vomiting or for	ever > 100.5 F trace give water, re-test keto	nes & bG in 2 hrs or_	hrs			
SC/IM Give PRN; unconsciou	SC/IM s. unresponsive.	seizure, or in			➤ If ketones	are moderate or large, give w	ater, Call parent and	Endocrinolo	gist □ NO GYM		
unknown. Tum onto let	t side to prevent	aspiration. If	more tha	n one option is	n one option is If ketones and vomiting, unable to take PO and MD not availa				ble, CALL 911		
chosen, school staff w	ill use ONE form	of available	glucagon	unless otherwise	☐ Give insul	in correction dose if > 2 hrs o	rhours since	last rapid act	ing insulin.		
directed.				SKILL LE	VEL (if not com	plete, will default to nurse-depend	entj				
Blood Glucose (bG)	Monitoring Skil			dministration Skill Lev		Independent Student	Self carry / Self-adr	ninister			
Nurse/adult must c			Nurse-	-Dependent Student: nu er medication	rse must	(MUST initial attestation).	I attest that the inde	pendent the prescrib	ed		
Student to check bo Student may check				vised student: student ca	alculates and	medication (excluding glu	ility to self-administer the prescribed ucagon) effectively during school,				
			self-admi	inisters, under adult sup	ervision	field trips and school spor	nsored events.		Provider Initials		
						e Part B for CGM reading					
	bG in school	(must match	times for	treatment and/or insulin	n) Break	fast □ Lunch □ Snack I □ Breakfast □ Lunch [ack before o	vm		
Hypoglycemia Check all boxes need	ed. Must includ	e at least one	e treatme	ent plan.			J SHACK LJ CIVE OII	1			
☐ For bG <	mg/dl give	gm rapid c	carbs at	☐ Breakfast ☐ Lunch	Breakfast □ Lunch □ Snack □ Gym □ PRN			☐ T2DM – no bG monitoring or insulin in school			
Repeat bG te	sting in 15 min o	ormir	n. If bG s	till <mg and="" bg="" carbs="" dl="" repeat="" retesting="" until="">mg/dl</mg>			g/dl				
☐ For bG <	_mg/dl give	gm rapid	carbs at	☐ Breakfast ☐ Lunc	inch ☐ Snack ☐ Gym ☐ PRN 15 gm ra			rapid carbs = 4			
Repeat bG testing in 15 min ormin. If bG still <mg and="" bg="" carbs="" dl="" repeat="" retesting="" until="">mg/dl</mg>											
Mid-Range Glycemia Insulin is given before food unless noted here Give insulin after Breakfast Lunch Snack Give Snack before gym if bG < mg/dl											
Mid-Range Glycemia	Insulin is a	iven before fo	ood unles	ss noted here	insulin after [□ Breakfast □ Lunch □	Snack ☐ Give Sna	ck before gy	ym if bG <mg dl<="" td=""></mg>		
Mid-Range Glycemia Hyperglycemia						☐ Breakfast ☐ Lunch ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐		ack before gy			
Hyperglycemia ☐ For bG	Insulin is g _mg/dl pre-gym	<i>iven before fo</i> 1, NO GYM	ood unles	ss noted here Give	insulin after 【	Breakfast □ Lunch □ For bG m					
Hyperglycemia ☐ For bG	Insulin is g _mg/dl pre-gym	<i>iven before fo</i> 1, NO GYM	ood unles		insulin after 【	☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin	Snack eter reading "High" u	se bG of 500	or mg/dl		
Hyperglycemia For bG For bG > Check bG or Sens	Insulin is g _mg/dl pre-gym _mg/dl PRN or Glucose (sG	iven before fon	correction	on dose if > 2 hrs or	insulin after [☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co	Snack eter reading "High" u prrection dose pre-me	se bG of 500	or mg/dl		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value	Insulin is g _mg/dl pre-gym _mg/dl PRN or Glucose (sG	iven before for	correction	on dose if > 2 hrs oremia if needed, and give	insulin after [☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co gm carb snack before dismis	Snack eter reading "High" u prrection dose pre-me sed	se bG of 500	or mg/dl		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value	Insulin is g _mg/dl pre-gym _mg/dl PRN or Glucose (sG	iven before for	correction	ss noted here	hrs. since	☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co gm carb snack before dismis bus/mass transit, parent to pic	Snack eter reading "High" u prrection dose pre-me sed	se bG of 500	or mg/dl		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value	Insulin is g _mg/dl pre-gym _mg/dl PRN or Glucose (sG	iven before for	correction	ss noted here Give on dose if > 2 hrs or emia if needed, and give plycemia if needed, and cive	hrs. since	☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co gm carb snack before dismis bus/mass transit, parent to pic	Snack eter reading "High" u prection dose pre-me sed k up from school.	se bG of 500	or mg/dl		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value	Insulin is g _mg/dl pre-gym _mg/dl PRN or Glucose (sG	iven before for	correction	ss noted here ☐ Give on dose if > 2 hrs or emia if needed, and give plycemia if needed, and or INSI Insulin Calculation M	hrs. since do not send on t ULIN ORDER	☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co gm carb snack before dismis bus/mass transit, parent to pic	Snack eter reading "High" u prection dose pre-me sed k up from school.	se bG of 500 al and carb	oor mg/dl coverage after meal (give number, not range)		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value	Insulin is g _mg/dl pre-gyn _mg/dl PRN, or Glucose (sG) s <mg< td=""><td>iven before fo</td><td>correction</td><td>on dose if > 2 hrs or emia if needed, and give_ glycemia if needed, and give_ INSI Insulin Calculation N Carb coverage ONI</td><td>hrs. since do not send on l ULIN ORDER lethod: LY at: Breakt</td><td>Breakfast Lunch For bG m For bG m a last rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic</td><td>Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time</td><td>se bG of 500 al and carb n Directions will be 7am to</td><td>oor mg/dl coverage after meal (give number, not range)</td></mg<>	iven before fo	correction	on dose if > 2 hrs or emia if needed, and give_ glycemia if needed, and give_ INSI Insulin Calculation N Carb coverage ONI	hrs. since do not send on l ULIN ORDER lethod: LY at: Breakt	Breakfast Lunch For bG m For bG m a last rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic	Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time	se bG of 500 al and carb n Directions will be 7am to	oor mg/dl coverage after meal (give number, not range)		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value	Insulin is g _mg/dl pre-gyn _mg/dl PRN, or Glucose (sG) s <mg< td=""><td>iven before fo</td><td>correction</td><td>on dose if > 2 hrs or</td><td>hrs. since hrs. since do not send on l ULIN ORDER Method: LY at: Break s correction do:</td><td>Breakfast Lunch For bG m For bG m I last rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic S fast Lunch Snack kfast Lunch Snack se when bG > Target AND</td><td>Snack eter reading "High" u prrection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG =</td><td>se bG of 500 al and carb n Directions will be 7am to</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified eto)</td></mg<>	iven before fo	correction	on dose if > 2 hrs or	hrs. since hrs. since do not send on l ULIN ORDER Method: LY at: Break s correction do:	Breakfast Lunch For bG m For bG m I last rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic S fast Lunch Snack kfast Lunch Snack se when bG > Target AND	Snack eter reading "High" u prrection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG =	se bG of 500 al and carb n Directions will be 7am to	or mg/dl coverage after meal (give number, not range) 4pm if not specified eto)		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG; s <m; <<="" es="" td=""><td>iven before for, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i</td><td>correction</td><td>on dose if > 2 hrs oremia if needed, and give_glycemia if needed, and give_glycemia if needed, and constant in the consta</td><td>hrs. since hrs. since do not send on le ULIN ORDER lethod: LY at: Break s correction doors since last rai</td><td>Breakfast Lunch For bG m For bG m Flast rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic S fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at</td><td>Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time</td><td>se bG of 500 al and carb n Directions will be 7am to</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified eto)</td></m;>	iven before for, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i	correction	on dose if > 2 hrs oremia if needed, and give_glycemia if needed, and give_glycemia if needed, and constant in the consta	hrs. since hrs. since do not send on le ULIN ORDER lethod: LY at: Break s correction doors since last rai	Breakfast Lunch For bG m For bG m Flast rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic S fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at	Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time	se bG of 500 al and carb n Directions will be 7am to	or mg/dl coverage after meal (give number, not range) 4pm if not specified eto)		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value Insulin Name*	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG; s <m; <<="" es="" td=""><td>iven before for, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i</td><td>correction</td><td>and dose if > 2 hrs oremia if needed, and give_glycemia if needed, and give_glycemia if needed, and color loss of the color loss of</td><td>hrs. since hrs. since do not send on le ULIN ORDER lethod: LY at: Break s correction do rs since last ra Lunch</td><td>Breakfast Lunch For bG m For bG m Flast rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic S fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at</td><td>Snack eter reading "High" u eter reading "Hi</td><td>n Directions will be 7am to _mg/dl (tim</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto)</td></m;>	iven before for, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i	correction	and dose if > 2 hrs oremia if needed, and give_glycemia if needed, and give_glycemia if needed, and color loss of the color loss of	hrs. since hrs. since do not send on le ULIN ORDER lethod: LY at: Break s correction do rs since last ra Lunch	Breakfast Lunch For bG m For bG m Flast rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic S fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at	Snack eter reading "High" u eter reading "Hi	n Directions will be 7am to _mg/dl (tim	or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto)		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <m; <m;="" c="" humalog<="" q="" s="" td="" with=""><td>iven before for, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i</td><td>correctionissal</td><td>and dose if > 2 hrs oremia if needed, and give_glycemia if needed, and give_glycemia if needed, and color loss of the color loss of</td><td>hrs. since hrs. since do not send on t ULIN ORDER Method: LY at: Break s correction do: rs since last rai Lunch Snoulated using:</td><td>☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co gm carb snack before dismis bus/mass transit, parent to pic S fast ☐ Lunch ☐ Snack kfast ☐ Lunch ☐ Snack se when bG > Target AND pid acting insulin at lack</td><td>Snack eter reading "High" u prrection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit</td><td>n Directions will be 7am tomg/dl (timmg/dl (timy Factor (IS)</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto)</td></m;>	iven before for, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i	correctionissal	and dose if > 2 hrs oremia if needed, and give_glycemia if needed, and give_glycemia if needed, and color loss of the color loss of	hrs. since hrs. since do not send on t ULIN ORDER Method: LY at: Break s correction do: rs since last rai Lunch Snoulated using:	☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co gm carb snack before dismis bus/mass transit, parent to pic S fast ☐ Lunch ☐ Snack kfast ☐ Lunch ☐ Snack se when bG > Target AND pid acting insulin at lack	Snack eter reading "High" u prrection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit	n Directions will be 7am tomg/dl (timmg/dl (timy Factor (IS)	or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto)		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value No Insulin Name* No Insulin in school Syringe/Pen	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <mg <<="" <mg="" ss="" td=""><td>iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack</td><td>correctionissal hypoglyce for hypog</td><td>andose if > 2 hrs oremia if needed, and giveelycemia if needed, and giveelycemia if needed, and constant if need</td><td>hrs. since hrs. since do not send on t ULIN ORDER Method: LY at: Break s correction do: rs since last raj Lunch Sn culated using:</td><td>☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co gm carb snack before dismis bus/mass transit, parent to pic S fast ☐ Lunch ☐ Snack kfast ☐ Lunch ☐ Snack se when bG > Target AND pid acting insulin at lack</td><td>Snack eter reading "High" u eter reading "Hi</td><td>n Directions will be 7am tomg/dl (timmg/dl (timy Factor (IS)</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto)</td></mg>	iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack	correctionissal hypoglyce for hypog	andose if > 2 hrs oremia if needed, and giveelycemia if needed, and giveelycemia if needed, and constant if need	hrs. since hrs. since do not send on t ULIN ORDER Method: LY at: Break s correction do: rs since last raj Lunch Sn culated using:	☐ Breakfast ☐ Lunch ☐ For bG m e last rapid acting insulin ☐ Give co gm carb snack before dismis bus/mass transit, parent to pic S fast ☐ Lunch ☐ Snack kfast ☐ Lunch ☐ Snack se when bG > Target AND pid acting insulin at lack	Snack eter reading "High" u eter reading "Hi	n Directions will be 7am tomg/dl (timmg/dl (timy Factor (IS)	or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto)		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <mg <<="" <mg="" ss="" td=""><td>iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack</td><td>correctionissal hypoglyce for hypog</td><td>andose if > 2 hrs or</td><td>hrs. since hrs. since do not send on le ULIN ORDER lethod: LY at: Break s correction do rs since last rap Lunch Since last rap Lun</td><td>Breakfast □ Lunch □ For bG m For bG m I last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pic IS fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract</td><td>Snack eter reading "High" u prrection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases</td><td>n Directions will be 7am to _mg/dl (timmg/dl (timmg/ds (timmg b G by</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dl</td></mg>	iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack	correctionissal hypoglyce for hypog	andose if > 2 hrs or	hrs. since hrs. since do not send on le ULIN ORDER lethod: LY at: Break s correction do rs since last rap Lunch Since last rap Lun	Breakfast □ Lunch □ For bG m For bG m I last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pic IS fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract	Snack eter reading "High" u prrection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases	n Directions will be 7am to _mg/dl (timmg/dl (timmg/ds (timmg b G by	or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dl		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen S Pump (Brand)	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <mg <<="" <mg="" ss="" td=""><td>iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack</td><td>correctionissal hypoglyce for hypog</td><td>an dose if > 2 hrs or</td><td>hrs. since hrs. Break s correction do rs since last rap Lunch</td><td>Breakfast □ Lunch □ For bG m For bG m I last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pic IS fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract</td><td>Snack eter reading "High" u eter reading "Hi</td><td>n Directions will be 7am tomg/dl (timmg/dl (tim y Factor (IS) bG by to</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dl</td></mg>	iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack	correctionissal hypoglyce for hypog	an dose if > 2 hrs or	hrs. since hrs. Break s correction do rs since last rap Lunch	Breakfast □ Lunch □ For bG m For bG m I last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pic IS fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract	Snack eter reading "High" u eter reading "Hi	n Directions will be 7am tomg/dl (timmg/dl (tim y Factor (IS) bG by to	or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dl		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen Pump (Brand) For Pumps:	Insulin is g mg/dl pre-gym mg/dl PRN, or Glucose (sG) s <m <="" humalog="" i<="" og="" ss="" td="" with=""><td>iven before for n, NO GYM Give insulin before dism g/dl treat for h, mg/dl treat in mg/dl treat in at Snack pen suggesting in the suggesting in at Snack</td><td>correctionissal hypoglyce for hypog</td><td>an dose if > 2 hrs or</td><td>hrs. since hrs. since hrs. since do not send on I ULIN ORDER Tethod: LY at: Break sty at: Brea</td><td>Breakfast □ Lunch □ For bG m last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pic S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract alculation.</td><td>Snack eter reading "High" u prrection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases</td><td>n Directions will be 7am tomg/dl (timmg/dl (tim y Factor (IS) bG by to</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dl</td></m>	iven before for n, NO GYM Give insulin before dism g/dl treat for h, mg/dl treat in mg/dl treat in at Snack pen suggesting in the suggesting in at Snack	correctionissal hypoglyce for hypog	an dose if > 2 hrs or	hrs. since hrs. since hrs. since do not send on I ULIN ORDER Tethod: LY at: Break sty at: Brea	Breakfast □ Lunch □ For bG m last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pic S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract alculation.	Snack eter reading "High" u prrection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases	n Directions will be 7am tomg/dl (timmg/dl (tim y Factor (IS) bG by to	or mg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dl		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen Pump (Brand) For Pumps: Student on FDA a	Insulin is g mg/dl pre-gym mg/dl PRN, or Glucose (sG) s <m <="" humalog="" i<="" og="" ss="" td="" with=""><td>iven before for n, NO GYM Give insulin before dism g/dl treat for h, mg/dl treat for h, and sales and sales are suggestive pen suggestive closed loop</td><td>correctionissal hypoglyce for hypog</td><td>an dose if > 2 hrs or</td><td>hrs. since hrs. since do not send on t ULIN ORDER Nethod: LY at: Break MLY at: Break MLY at: Break since last rap Lunch Since last rap Lunch Sinther Orders) Part B) mediately follo m lunch carb catructions: mmendations fo</td><td>Breakfast □ Lunch □ For bG m For bG m I last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pic IS fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract</td><td>Snack eter reading "High" u eter reading "Hi</td><td>n Directions will be 7am to mg/dl (tim mg/dl (tim y Factor (IS) bG by to bG by</td><td>ormg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl</td></m>	iven before for n, NO GYM Give insulin before dism g/dl treat for h, mg/dl treat for h, and sales and sales are suggestive pen suggestive closed loop	correctionissal hypoglyce for hypog	an dose if > 2 hrs or	hrs. since hrs. since do not send on t ULIN ORDER Nethod: LY at: Break MLY at: Break MLY at: Break since last rap Lunch Since last rap Lunch Sinther Orders) Part B) mediately follo m lunch carb catructions: mmendations fo	Breakfast □ Lunch □ For bG m For bG m I last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pic IS fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract	Snack eter reading "High" u eter reading "Hi	n Directions will be 7am to mg/dl (tim mg/dl (tim y Factor (IS) bG by to bG by	ormg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen Pump (Brand) For Pumps: Student on FDA a pump-basal rate variance	Insulin is g mg/dl pre-gym mg/dl PRN, or Glucose (sG) s <m <="" es="" humalog="" i<="" og="" td="" with=""><td>iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat in at Snack pen suggestion closed loop</td><td>correctionissal hypoglyce for hypog</td><td>an dose if > 2 hrs or</td><td>hrs. since hrs. since do not send on t ULIN ORDER Nethod: LY at: Break MLY at: Break MLY at: Break se correction doe rs since last rap Lunch Sn culated using: bither Orders) Part B) mediately follo medi</td><td>Breakfast □ Lunch □ For bG m last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pid S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract alculation.</td><td>Snack eter reading "High" u eter reading "Hi</td><td>n Directions will be 7am to _mg/dl (tim _mg/dl (tim _mg/dl (tim _to_bg bytotototototo</td><td>ormg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl</td></m>	iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat in at Snack pen suggestion closed loop	correctionissal hypoglyce for hypog	an dose if > 2 hrs or	hrs. since hrs. since do not send on t ULIN ORDER Nethod: LY at: Break MLY at: Break MLY at: Break se correction doe rs since last rap Lunch Sn culated using: bither Orders) Part B) mediately follo medi	Breakfast □ Lunch □ For bG m last rapid acting insulin □ Give co gm carb snack before dismis bus/mass transit, parent to pid S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract alculation.	Snack eter reading "High" u eter reading "Hi	n Directions will be 7am to _mg/dl (tim _mg/dl (tim _mg/dl (tim _to_bg bytotototototo	ormg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in schoo Delivery Method Syringe/Pen Pump (Brand) For Pumps: Student on FDA a pump-basal rate vari Suspend/disconn	Insulin is g mg/dl pre-gym mg/dl PRN, or Glucose (sG) s <m <="" es="" humalog="" i<="" og="" td="" with=""><td>iven before for n, NO GYM Give insulin of before dism g/dl treat for h, mg/dl treat in mg/dl tre</td><td>correctionissal appropriate for hypograms ions</td><td>an dose if > 2 hrs or</td><td>hrs. since hrs. since do not send on t ULIN ORDER Method: LY at: Break LY at: Break s correction do rs since last ra Lunch Sn culated using: bither Orders) Part B) mediately follow m lunch carb co tructions: mmendations fo ons, will round of g/dl that has no</td><td>Breakfast Lunch For bG m For bG m Plast rapid acting insulin Give composed grant snack before dismiss bus/mass transit, parent to pices Fast Lunch Snack before dismiss bus/mass transit, parent to pices Fast Lunch Snack before dismiss grant for pices Fast Lunch Snack before dismiss grant for pices Fast Lunch Snack before dismission grant for grant</td><td>Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R</td><td>n Directions will be 7am tomg/dl (timmg/dl) tobG by totoatio (I:C):</td><td>coverage after meal (give number, not range) (4pm if not specified eto) eto) F1:mg/dlng/dl</td></m>	iven before for n, NO GYM Give insulin of before dism g/dl treat for h, mg/dl treat in mg/dl tre	correctionissal appropriate for hypograms ions	an dose if > 2 hrs or	hrs. since hrs. since do not send on t ULIN ORDER Method: LY at: Break LY at: Break s correction do rs since last ra Lunch Sn culated using: bither Orders) Part B) mediately follow m lunch carb co tructions: mmendations fo ons, will round of g/dl that has no	Breakfast Lunch For bG m For bG m Plast rapid acting insulin Give composed grant snack before dismiss bus/mass transit, parent to pices Fast Lunch Snack before dismiss bus/mass transit, parent to pices Fast Lunch Snack before dismiss grant for pices Fast Lunch Snack before dismiss grant for pices Fast Lunch Snack before dismission grant for grant	Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R	n Directions will be 7am tomg/dl (timmg/dl) tobG by totoatio (I:C):	coverage after meal (give number, not range) (4pm if not specified eto) eto) F1:mg/dlng/dl		
Hyperglycemia For bG For bG > Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen S Pump (Brand) For Pumps: Student on FDA a pump-basal rate variance Suspend/disconn	Insulin is g mg/dl pre-gym mg/dl PRN, or Glucose (sG) s <m humalog="" i<="" og="" td="" with=""><td>iven before for n, NO GYM Give insulin of before dism g/dl treat for h, mg/dl treat in mg/dl tre</td><td>correctionissal appropriate for hypograms ions</td><td>ss noted here Give on dose if > 2 hrs or</td><td>hrs. since hrs. since do not send on I ULIN ORDER Nethod: LY at: Break Scorrection do: rs since last ra Lunch Sn culated using: part B) mediately follor mediately follor mediations: nmendations fo ons, will round of gdid that has no der pump failure: SUSI</td><td>For bG m For bG m For bG m For bG m For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at tack □ ISF or □ Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased inhours re and notify parents. PEND pump, give rapid</td><td>Snack eter reading "High" u eter reading "Hi</td><td>n Directions will be 7am tomg/dl (timmg/dl) tobG by totoatio (I:C):</td><td>coverage after meal (give number, not range) (4pm if not specified eto) eto) F1:mg/dlng/dl</td></m>	iven before for n, NO GYM Give insulin of before dism g/dl treat for h, mg/dl treat in mg/dl tre	correctionissal appropriate for hypograms ions	ss noted here Give on dose if > 2 hrs or	hrs. since hrs. since do not send on I ULIN ORDER Nethod: LY at: Break Scorrection do: rs since last ra Lunch Sn culated using: part B) mediately follor mediately follor mediations: nmendations fo ons, will round of gdid that has no der pump failure: SUSI	For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at tack □ ISF or □ Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased inhours re and notify parents. PEND pump, give rapid	Snack eter reading "High" u eter reading "Hi	n Directions will be 7am tomg/dl (timmg/dl) tobG by totoatio (I:C):	coverage after meal (give number, not range) (4pm if not specified eto) eto) F1:mg/dlng/dl		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in schoo Delivery Method Syringe/Pen Pump (Brand) For Pumps: Student on FDA a pump-basal rate vari Suspend/disconn	Insulin is g mg/dl pre-gym mg/dl PRN, or Glucose (sG) s <m <="" es="" humalog="" i<="" og="" td="" with=""><td>iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat in mg/dl treat i</td><td>correctionissal appropriate for hypograms ions</td><td>ss noted here Give on dose if > 2 hrs or</td><td>hrs. since hrs. since do not send on I ULIN ORDER Nethod: LY at: Break Scorrection do: rs since last ra Lunch Sn culated using: Other Orders) Part B) Immediately follor mediately follor mediately follor mediations fo ons, will round of gridl that has no der pump faillure: SUSI ge or pen, and</td><td>For bG m For bG m For bG m For bG m For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S Fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at tack ISF or Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased in hours re and notify parents. PEND pump, give rapid notify parents.</td><td>Snack eter reading "High" u eter reading send in the s</td><td>n Directions will be 7am to mg/dl (tim mg/dl (tim y Factor (IS bG by to bG by to atio (I:C):</td><td>ormg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl</td></m>	iven before for n, NO GYM Give insulin before dism g/dl treat for h mg/dl treat in mg/dl treat i	correctionissal appropriate for hypograms ions	ss noted here Give on dose if > 2 hrs or	hrs. since hrs. since do not send on I ULIN ORDER Nethod: LY at: Break Scorrection do: rs since last ra Lunch Sn culated using: Other Orders) Part B) Immediately follor mediately follor mediately follor mediations fo ons, will round of gridl that has no der pump faillure: SUSI ge or pen, and	For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S Fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at tack ISF or Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased in hours re and notify parents. PEND pump, give rapid notify parents.	Snack eter reading "High" u eter reading send in the s	n Directions will be 7am to mg/dl (tim mg/dl (tim y Factor (IS bG by to bG by to atio (I:C):	ormg/dl coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen Student on FDA a pump-basal rate variable suspend/disconn Suspend pump for to treatment for	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <mg <="" humalog="" l<="" og="" ss="" td="" with=""><td>iven before for n, NO GYM Give insulin Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack pen suggesti closed loop m a not respond</td><td>correctionissal hypoglyce for hypoglyce for</td><td>ss noted here Give on dose if > 2 hrs or</td><td>hrs. since hrs. since do not send on the send of the s</td><td>For bG m For bG m For bG m For bG m For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at tack □ ISF or □ Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased inhours re and notify parents. PEND pump, give rapid</td><td>Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R Bkfast OR time 1 unit per</td><td>n Directions al and carb in Directions will be 7am tomg/dl (timmg/dl (</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified e to) e to) F1: mg/dl mg/dl) arbs</td></mg>	iven before for n, NO GYM Give insulin Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack pen suggesti closed loop m a not respond	correctionissal hypoglyce for	ss noted here Give on dose if > 2 hrs or	hrs. since hrs. since do not send on the send of the s	For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at tack □ ISF or □ Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased inhours re and notify parents. PEND pump, give rapid	Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R Bkfast OR time 1 unit per	n Directions al and carb in Directions will be 7am tomg/dl (timmg/dl (or mg/dl coverage after meal (give number, not range) 4pm if not specified e to) e to) F1: mg/dl mg/dl) arbs		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen Suspend/disconn Suspend/disconn Suspend pump for treatment for Activity Mode (HStart minutes gduration (DEFAULT)	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <mg <="" humalog="" i<="" og="" ss="" td="" with=""><td>iven before for n, NO GYM Give insulin Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack pen suggesti closed loop m a not respond</td><td>correctionissal hypoglyce for hypoglyce for</td><td>ss noted here Give on dose if > 2 hrs or</td><td>hrs. since hrs. since do not send on the send of the s</td><td>For bG m For bG m For bG m For bG m For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S Fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at tack ISF or Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased in hours re and notify parents. PEND pump, give rapid notify parents.</td><td>Snack eter reading "High" u eter reading send in the s</td><td>n Directions al and carb in Directions will be 7am tomg/dl (timmg/dl (</td><td>or mg/dl coverage after meal (give number, not range) 4pm if not specified e to) e to) F1: mg/dl mg/dl) arbs</td></mg>	iven before for n, NO GYM Give insulin Give insulin before dism g/dl treat for h mg/dl treat i //Admelog n at Snack pen suggesti closed loop m a not respond	correctionissal hypoglyce for	ss noted here Give on dose if > 2 hrs or	hrs. since hrs. since do not send on the send of the s	For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S Fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at tack ISF or Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased in hours re and notify parents. PEND pump, give rapid notify parents.	Snack eter reading "High" u eter reading send in the s	n Directions al and carb in Directions will be 7am tomg/dl (timmg/dl (or mg/dl coverage after meal (give number, not range) 4pm if not specified e to) e to) F1: mg/dl mg/dl) arbs		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen Suspend/disconn Suspend/disconn Suspend pump for to treatment for Activity Mode (HStart minutes processed) minutes are sure following exercise)	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <mg <="" humalog="" i<="" og="" ss="" td="" with=""><td>iven before for n, NO GYM Give insulin of the dismonth of the</td><td>correctionissal hypoglyce for hypoglyce for</td><td>an dose if > 2 hrs or</td><td>hrs. since hrs. since hrs. since do not send on t ULIN ORDER Method: LY at: Break struction do: rs since last ra; Lunch Soulated using: Other Orders) Part B) Immediately follow method to tructions: ons, will round of gold that has no der pump failure; SUSI ge or pen, and re, only give corresultin</td><td>For bG m For bG m For bG m For bG m For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S Fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at tack ISF or Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased in hours re and notify parents. PEND pump, give rapid notify parents.</td><td>Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R Bkfast OR time 1 unit per</td><td>n Directions al and carb in Directions will be 7am to mg/dl (tim mg/dl (tim mg/dl (tim y Factor (IS) bG by to bG by to atio (I:C):</td><td>coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl</td></mg>	iven before for n, NO GYM Give insulin of the dismonth of the	correctionissal hypoglyce for	an dose if > 2 hrs or	hrs. since hrs. since hrs. since do not send on t ULIN ORDER Method: LY at: Break struction do: rs since last ra; Lunch Soulated using: Other Orders) Part B) Immediately follow method to tructions: ons, will round of gold that has no der pump failure; SUSI ge or pen, and re, only give corresultin	For bG m Give co Gm carb snack before dismis bus/mass transit, parent to pic S Fast Lunch Snack kfast Lunch Snack se when bG > Target AND pid acting insulin at tack ISF or Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) tt decreased in hours re and notify parents. PEND pump, give rapid notify parents.	Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R Bkfast OR time 1 unit per	n Directions al and carb in Directions will be 7am to mg/dl (tim mg/dl (tim mg/dl (tim y Factor (IS) bG by to bG by to atio (I:C):	coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Pump (Brand) For Pumps: Student on FDA a pump-basal rate vari Suspend/disconn Suspend pump fot to treatment for Activity Mode (HStart minutes guration (DEFAULT following exercise) Carb Coverage:	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <mg <="" humalog="" i<="" og="" ss="" td="" with=""><td>iven before for n, NO GYM Give insulin Give insulin Defore dism g/dl treat in g/dl treat in g/Admelog n at Snack pen suggestin closed loop m a not respond for min g, and 2 hrs ction Dose usin</td><td>correctionissal hypoglyce for hypoglyce for</td><td>ss noted here</td><td>hrs. since hrs. since do not send on I ULIN ORDER Method: LY at: Break LY at: Break scorrection do rs since last ra Lunch Sn culated using: bither Orders) Part B) mediately follor m lunch carb ca fructions: mmendations fo ns, will round of gold that has no der pump failure; SUSI ge or pen, and e, only give corr sulin e to closest 0.5 un oesn't have ½ unit</td><td>For bG m le last rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) t decreased inhours re and notify parents. PEND pump, give rapid notify parents. ection dose if >hrs iit for syringe/pen, or nearest marks; unless otherwise</td><td>Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R Bkfast OR time 1 unit per Snack OR time 1 unit per 1 unit per 1 unit per 1 unit per</td><td>n Directions will be 7am to mg/dl (tim mg/dl (tim y Factor (IS bG by to bG by to atio (I:C):</td><td>coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dl) arbs oarbs</td></mg>	iven before for n, NO GYM Give insulin Give insulin Defore dism g/dl treat in g/dl treat in g/Admelog n at Snack pen suggestin closed loop m a not respond for min g, and 2 hrs ction Dose usin	correctionissal hypoglyce for	ss noted here	hrs. since hrs. since do not send on I ULIN ORDER Method: LY at: Break LY at: Break scorrection do rs since last ra Lunch Sn culated using: bither Orders) Part B) mediately follor m lunch carb ca fructions: mmendations fo ns, will round of gold that has no der pump failure; SUSI ge or pen, and e, only give corr sulin e to closest 0.5 un oesn't have ½ unit	For bG m le last rapid acting insulin Give co gm carb snack before dismis bus/mass transit, parent to pic S fast □ Lunch □ Snack kfast □ Lunch □ Snack se when bG > Target AND pid acting insulin at lack □ ISF or □ Sliding Scale wing lunch, subtract alculation. In bolus dose (if not using down to nearest 0.1 unit) t decreased inhours re and notify parents. PEND pump, give rapid notify parents. ection dose if >hrs iit for syringe/pen, or nearest marks; unless otherwise	Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R Bkfast OR time 1 unit per Snack OR time 1 unit per 1 unit per 1 unit per 1 unit per	n Directions will be 7am to mg/dl (tim mg/dl (tim y Factor (IS bG by to bG by to atio (I:C):	coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dl) arbs oarbs		
Hyperglycemia For bG Check bG or Sens For sG or bG value For sG or bG value Insulin Name* *May substitute Novol No Insulin in school Delivery Method Syringe/Pen Suspend/disconn Suspend/disconn Suspend pump for to treatment for Activity Mode (HStart minutes processed) minutes are sure following exercise)	Insulin is g _mg/dl pre-gym _mg/dl PRN, or Glucose (sG) s <mg humalog="" l<="" og="" s="" td="" with=""><td>iven before for n, NO GYM Give insulin of the dismonth of the</td><td>correctionissal hypoglyce for hypoglyce for</td><td>ss noted here</td><td>hrs. since hrs. since hrs. since do not send on I ULIN ORDER Nethod: LY at: Break Scorrection do: rs since last ral Lunch Sn culated using: Other Orders) Part B) Immediately follor mediately follor mediately</td><td>For bG m For bG m For bG m For bG m Give co Give co</td><td>Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R Bkfast OR time 1 unit per Snack OR time</td><td>n Directions al and carb in Directions will be 7am to mg/dl (time mg/dl (time y Factor (IS) bG by to bG by to atio (I:C): gms co to gms co to</td><td>coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl arbs</td></mg>	iven before for n, NO GYM Give insulin of the dismonth of the	correctionissal hypoglyce for	ss noted here	hrs. since hrs. since hrs. since do not send on I ULIN ORDER Nethod: LY at: Break Scorrection do: rs since last ral Lunch Sn culated using: Other Orders) Part B) Immediately follor mediately	For bG m For bG m For bG m For bG m Give co	Snack eter reading "High" u prection dose pre-me sed k up from school. Insulin Calculatio If only one given, time Target bG = Insulin Sensitivit 1 unit decreases (time 1 unit decreases (time Insulin to Carb R Bkfast OR time 1 unit per Snack OR time	n Directions al and carb in Directions will be 7am to mg/dl (time mg/dl (time y Factor (IS) bG by to bG by to atio (I:C): gms co to gms co to	coverage after meal (give number, not range) 4pm if not specified eto) eto) F1:mg/dlmg/dl arbs		



Diabetes Medication Administration Form [Part B]

Provider Medication Order Form | School Year 2023-24 Please fax all DMAFs to 347-396-8932/8945

DUE: June 1st. Forms submitted after June 1st may delay processing for new school year.

Student Last Name First Name					OSIS#					
CONTINUOUS GLUCOSE MONITORING (CGM) ORDERS [Please see 'Provider Guidelines for DMAF Completion']										
		I's used to repla	ace finger sticl	k bG readings, nodel of the CG	only devices FD M in use.	A approved for use and age Model of CGM:				rer's
	for insulin dosing: nsor does not sho			hen: the symp	toms don't mate	h the CGM readings; if there			1500	adings
□ CGM to be used for insulin dosing and monitoring - must be FDA approved for use and age <u>sG Monitoring</u> Specify times to check sensor reading □ Breakfast □ Lunch □ Snack □ Gym □ PRN [if none checked, will use bG monitoring times] For sG <70mg/dL check bG and follow orders on DMAF, unless otherwise ordered below. Use CGM grid below OR □ See attached CGM instruction										
CGM reading		Arrows Action □ use < 80 mg/dl instead of < 70 mg/dl for grid action plan								
sG < 60 mg/dl	, , , , , , , , , , , , , , , , , , , ,									
sG 60-70 mg/dl		and 1, 11,	or →			r bG hypoglycemia plan; Rec				
sG 60-70 mg/dl	6 60-70 mg/dl and ↑ , ↑↑, or >			If still <	70 mg/dl check l		emia plan; if no	ot symptomatic, re	echeck in 15-20) minutes.
sG >70 mg/dl		Any arrows	S			s for insulin dosing				
sG ≤ 120 mg/dl recess	G ≤ 120 mg/dl pre-gym or and ↓, ↓↓			i gms uncovered lculation	d carbs. If gym or recess is im	mediately after	· lunch, subtract 1	5 gms of carbs	s from lunch	
sG ≥ 250		Any arrows	3			s for treatment and insulin do	sing			
☐ For student u	using CGM, wait 2	hours after me	al before testi	ng ketones with	n hyperglycemia					
				PARENTAL	INPUT INTO	INSULIN DOSING				
Parent(s)/Guardian(s) (give name), may provide the nurse with information relevant to insulin dosing, including dosing recommendations. Taking the parent's input into account, the nurse will determine the insulin dose within the range ordered by the health care practitioner and in keeping with nursing judgment.										
4		-11-1-1			ase select ONE	2. Nurse may adju	ust calculated o	ose un hy	% or down by	%
	urse may adjust c arental input and			o to ur	nts based	of the prescribed				
MUST COMPLETE: Health care practitioner can be reached for urgent dosing orders at: () If the parent requests a similar adjustment for > 2 days in a row, the nurse will contact the health care practitioner to see if the school orders need to be revised.									a similar	
		SLIDING S	CALE			T	OPTION	AL ORDERS		
dose will be give	p ranges (e.g. ent en. Use pre-treatm	ter 0-100, 101-	200, etc.). If r			☐ Round insulin dosing to ☐ Round insulin dosing to half unit syringe/pen).	nearest whole	unit: 0,51-1,50u		101
☐ Lunch ☐ Snack	bG	Units O Insulin	ther Time	bG	Units Insulin	☐ Use sliding scale for co	_			
☐ Breakfast ☐ Correction	Zero -		Lunch Snack	Zero -		units for lunch;units for snack;units for breakfast				
Dose	386		Breakfast	*		(sliding scale must be marked as correction dose only)				
			Correction Dose			☐ Long-acting insulin gi	iven in school	– Insulin Name:		
	4					Dose:unit	s Time	or	☐ Lunch	
OTHER ORDI	ERS					HOME MEDICATIONS		□ None		
					_	Medication Insulin	Dose	Frequency	Time	Route
						TISUIII				
						Other				
						+				
					-				1	
				AD	DITIONAL IN	FORMATION				
Is the child us	ing altered or non-	FDA approved				te that New York State Educa tk up orders on DMAF Part A For		bit nurses from m	anaging non-F	DA devices.
			ng this form,	I certify that I	have discusse	d these orders with the par				
Health Care Practition	Health Care Practitioner LAST FIRST SIGNATURE DATE									
PLEASE PRINT	check one	MD DC		□ PA			111			
Address STREET			C	CITY/STATE		ZIP	Email			
NPI# or NYS License # (Required) Tel					Fax		CDC & AAP recommend annual seasonal influenza vaccination for all children diagnosed with diabetes.			

Office of School Health

Diabetes Medication Administration Form

Provider Medication Order Form | School Year 2023-24 Please fax all DMAFs to 347-396-8932/8945

DUE: June 1st. Forms submitted after June 1st may delay processing for new school year.

PARENTS AND GUARDIANS: READ, COMPLETE, AND SIGN. BY SIGNING BELOW, I AGREE TO THE FOLLOWING:

- 1. I consent to the nurse/school based health center (SBHC) provider giving my child's prescribed medicine, and the nurse/trained staff/SBHC provider checking their blood sugar and treating their low blood sugar based on the directions and skill level determined by my child's health care practitioner. These actions may be performed on school grounds or during school trips.
- 2. I also consent to any equipment needed for my child's medicine being stored and used at school.

3. I understand that:

- I must give the school nurse/SBHC provider my child's medicine, snacks, equipment, and supplies and must replace such medicine, snacks, equipment and supplies as needed. The Office of School Health (OSH) recommends the use of safety lancets and other safety needle devices and supplies to check my child's blood sugar levels and give insulin.
- All prescription and "over-the-counter" medicine I give the school must be new, unopened, and in the original bottle or box. I will provide the school with current, unexpired medicine for my child's use during school days.
 - Prescription medicine must have the original pharmacy label on the box or bottle. Label must include: 1) my child's name, 2) pharmacy name and phone number, 3) my child's health care practitioner's name, 4) date, 5) number of refills, 6) name of medicine, 7) dosage, 8) when to take the medicine, 9) how to take the medicine and 10) any other directions.
- I must **immediately** tell the school nurse/SBHC provider about any change in my child's medicine or the health care practitioner's instructions.
- OSH and its agents involved in providing the above health service(s) to my child are relying on the accuracy of the information in this form.
- By signing this Medication Administration Form (MAF), I authorize OSH to provide diabetes-related health services to my child. Theseservices may include but are not limited to a clinical assessment or a physical exam by an OSH health care practitioner or nurse.
- The medication order in this MAF expires at the end of my child's school year, which may include the summer session, or when I give the school nurse/SBHC provider a new MAF (whichever is earlier). When this medication order expires, I will give my child's school nurse/SBHC provider a new MAF written by my child's health care practitioner.
- OSH and the Department of Education (DOE) make sure that my child can safely test their blood sugar.
- This form represents my consent and request for the diabetes services described on this form. It is not an agreement by OSH to provide
 the requested services. If OSH decides to provide these services, my child may also need a Section 504 Accommodation Plan.
 This plan will be completed by the school.
- For the purposes of providing care or treatment for my child, OSH may obtain any other information they think is needed about my child's medical condition, medication or treatment. OSH may obtain this information from any health care practitioner, nurse, or pharmacist who has given my child health services.

OSH Parent Hotline for questions about the Diabetes Medication Administration Form (DMAF): 718-310-2496

FOR SELF-ADMINISTRATION OF MEDICINE (INDEPENDENT STUDENTS ONLY)

- I certify/confirm that my child has been fully trained and can take medicine on their own. I consent to my child carrying, storing and giving them the medicine prescribed on this form in school. I am responsible for giving my child this medicine in bottles or boxes as described above. I am also responsible for monitoring my child's medication use, and for all results of my child's use of this medicine in school. The school nurse or SBHC providers will confirm my child's ability to carry and give them medicine. I also agree to give the school "back up" medicine in a clearly labeled box or bottle.
- I consent to the school nurse or trained school staff giving my child Glucagon if prescribed by their health care provider if my child is temporarily unable to carry and take medicine.

NOTE: It is preferred that you send medication and equipment for your child on a school trip day and for off-site school activities.

Parent / Guardian'	Borough	Data signed	District		
Parent / Guardian'	's Signature for Parts A & B	Date signed			
	Parent / Guardian's Signature for Parts A & B Date		Date signed		
	Parent /Guardian's Email				
Home Tel No.		Cell Phone No.			
Relationship to St	Relationship to Student		Contact Tel No.		
		Home Tel No.	Home Tel No. Cell Phone No.		



Diabetes Medication Administration Form

Provider Medication Order Form | School Year 2023-24 Please fax all DMAFs to 347-396-8932/8945

For Office of School Health (OSH) Use Only

OSIS Number:							
Received by: Name	Date://						
Reviewed by: Name	Date:						
□504 □IEP □Other	Referred to School 504 Coordinator ☐ Yes ☐ No						
Services provided by: Nurse/NP School Based Health Center	OSH Public Health Advisor (for supervised students only)						
Signature and Title (RN OR SMD):							
Date School Notified & Form Sent to DOE Liaison							
Revisions as per OSH contact with prescribing health care practitioner Clarified Modified							
Notes							